S41B - 0997 Seismic Energy Partitioning Inferred from Pseudotachylytebearing Faults (Gole Larghe Fault, Adamello batholith, Italy) AGU FALL MEETING Lidia Pittarello* Giulio Di Toro* Andrea Bizzarri** Jafar Hadizadeh*** Giorgio Pennacchioni*

lidia.pittarello@unipd.it giulio.ditoro@unipd.it bizzarri@bo.ingv.it *Dipartimento di Geologia, Paleontologia e Geofisica, Università di Padova, Italy **Istituto Nazionale di Geofisica e Vulcanologia, Bologna, Italy

hadizadeh@louisville.edu giorgio.pennacchioni@unipd.it ***Department of Geography and Geosciences, University of Louisville, Kentucky USA

Aim of this study

Partitioning of the earthquake energy between fracture energy E_c (energy required to create new rupture surface in the slip zone and a damage zone in the wall rock) and frictional heat E_µ determines the features of the rupture propagation and the mechanical behavior of a seismic fault. The E_{c}/E_{μ} ratio cannot be inferred from seismological investigations. We propose to use the cataclastic microstructures associated with pseudotachylyte (solidified friction melt produced during coseismic slip) to constrain the E_{c}/E_{μ} ratio.

Methods

1. We selected a pseudotachylyte-bearing fault, that records one single seismic rupture, from an exhumed fault exposed in the Adamello batholith (Gole Larghe Fault zone, Italy, Pan.1). **2.** We estimated E_{μ} by energy balance calculations (Pan.2). **3.** We estimated *E*_c by: 3a. SEM and FE-SEM image analysis of fragmented plagioclase survivor clasts within the pseudotachylyte and fracture patterns in the host rock. 3b. Clast Size Distribution (CSD) and fracture density by computer-aided image analysis. E_{c} , then, was determined by multiplying the seismically created new fracture surfaces for the specific surface energy (γ) of the rock-forming minerals (Pan.4).

Pan.1-Geological Setting



Geological map of the Adamello batholith (in gray). Location of the map (black box) is shown in the inset. The yellow star marks the studied area.

The Gole Larghe Fault is an E-W trending dextral strike-slip fault which crosscuts the Adamello tonalitic batholith (Italian Southern Alps). Ambient conditions during seismic faulting were 250<T<300 ^oC and 250<P<300 MPa (Ref. 1).

Results & Conclusions

The above estimates yield $E_{\mu} = 23.3 \text{ MJ m}^{-2}$ and E_{σ} in the range of 0.110-0.500 MJ m⁻².

We conclude that, for this local seismic energy balance estimate, E_c is negligible compared to E_{μ} (Pan.5).

	IHζ		çř
Tonalite		Austroal	oine
Southalpine	<u> </u>	Faults	

faults Pseudotachylyte-bearing exploited preexisting joints. Some of these faults record one seismic rupture (Pan.3)

		10 cm
	tonalite	
pseudotachy	/lyte	

Pan.2-Estimate of E From Ref. 2: $E_{\rm H} = [(1-\phi) H + c_{\rm P} (T_{\rm m} - T_{\rm hr})] \rho 2W$ where ϕ = ratio clasts/pt matrix (0.2) (Fig.4a), H= latent heat of fusion (3.28 10^5 J kg⁻¹), **C**_b = the specific heat (1180 J kg⁻¹ K⁻¹), $T_m =$ initial friction melt temperature (1723 K) T_{hr} = host rock temperature (523 K), ρ = melt density (2350 kg m⁻³) and 2w = average pseudotachylyte thickness. For the 5.9 10⁻³ m thick fault shown in Pan.3:

 $E_{\rm H} = 23.3 \,\rm MJ \,m^{-2}$



of the pseudotachylyte.

• We determined the fragment distribution (CSD) within clasts by computer-aided image analysis on SEM and FE-SEM images (Fig.c-f). Two examples of CSD, measured from images at different magnifications, are shown in Fig.g. The CSD is not fractal over the whole range of measured sizes r $(0.05-100 \ \mu m)$. We identify three average "fractal" dimension: (i) D = 0,1 for $0.05 < r < 0.50 \mu m$, (ii) D = 0.85 for 0.50 < r < 1.00 μ m, (iii) D = 3.5 for 1.00 < r < 100 µm.





• We determined the fragment total surface per unit area (Sf) by assuming: (i) a spherical shape of the fragments and (ii) using the different D values for each grain size class. The Sf value is in the range 11000 and 50000.

We determined the fracture density in the host rock (Lf) on two orthogonal sections. The Lf values on the two sections are comparable. Whatever the assumption on the exact 3D geometry of fractures, the measured Lf values indicate that the host rock fracture surface is neglible compared to Sf.

Given $E_{g} = Sf * \gamma (Ref.3,4)$ and $\gamma = 10 J m^{-2} (Ref.5)$:

0.11< *E*_G <0.50 MJ m⁻²

Pan.5 - Conclusions

clasts

Our study yields a local estimate of E_{G}/E_{H} in the range 0.005-0.02. Assuming that the amount of energy radiated as seismic waves (E_s) represents the 0,1-10% of the total energy of an earthquake (Ref.6), we suggest the partitioning of E_s , E_c and E_{μ} reported in the pie-diagram below for this local context.

quartz

100µm WD 10.0mn

X70



References:

1. Di Toro, G., Pennacchioni, G., 2004. Superheated friction-induced melts in zoned pseudotachylytes within the Adamello tonalites (Italian Southern *Alps*) J. Struct. Geol. 26, 1783–1801;

2. Di Toro, G., et al., 2005. Can pseudotachylytes be used to infer earthquake source parameters? An example of limitations in the study of exhumed faults, Tectonophysics, 402, 3-20;

3. Chester, J., et al., 2005. Fracture surface energy of the Punchbowl fault, San Andreas system, Nature, 437, 133-136;

4. Wilson, B., et al., 2005. Particle size and energetics of gouge from earthquake rupture zones, Nature, 434, 749-752

5. Brace, W.F., Walsh, J.B., 1962. Some direct measurements of the surface energy of quartz and orthoclase, Am. Min. 47, 1111-1122.

6. McGarr, A., 1999. On relating apparent stress to the stress causing earthquake slip, Journal of Geophysical Research 104, 3003–3011.